




# Bifurcation Theory

## Lecture 2

a.y. 2013/14



# STABILITY AND BIFURCATION OF DYNAMICAL SYSTEMS

ANGELO LUONGO

## **Scope:**

- To remind basic notions of Dynamical Systems and Stability Theory;
- To introduce fundamentals of Bifurcation Theory;

## **Outline:**

1. General definitions
2. Fundamentals of Stability Theory
3. Fundamentals of Bifurcation Theory
4. Multiple bifurcations from a known path

# 1. GENERAL DEFINITIONS

We give general definitions for a  $N$ -dimensional autonomous systems.

- Equations of motion:

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t)), \quad \mathbf{x} \in \mathbb{R}^N$$

where  $\mathbf{x}$  are *state-variables*,  $\{\mathbf{x}\}$  the *state-space*, and  $\mathbf{F}$  the *vector field*.

- Orbits:

Let  $\mathbf{x}_s(t)$  be the solution to equations which satisfies prescribed initial conditions:

$$\begin{cases} \dot{\mathbf{x}}_s(t) = \mathbf{F}(\mathbf{x}_s(t)) \\ \mathbf{x}_s(0) = \mathbf{x}^0 \end{cases}$$

The set of all the values assumed by  $\mathbf{x}_s(t)$  for  $t > 0$  is called an *orbit* of the dynamical system. Geometrically, an orbit is a curve in the phase-space, originating from  $\mathbf{x}^0$ . The set of all orbits is the *phase-portrait* or *phase-flow*.

- Classifications of orbits:

Orbits are classified according to their time-behavior.

- *Equilibrium (or fixed-) point*: it is an orbit  $\mathbf{x}_S(t) =: \mathbf{x}_E$  independent of time (represented by a point in the phase-space);
- *Periodic orbit*: it is an orbit  $\mathbf{x}_S(t) =: \mathbf{x}_P(t)$  such that  $\mathbf{x}_P(t+T) = \mathbf{x}_P(t)$ , with  $T$  the period (it is a closed curve, called *cycle*);
- *Quasi-periodic orbit*: it is an orbit  $\mathbf{x}_S(t) =: \mathbf{x}_Q(t)$  such that, given an arbitrary small  $\varepsilon > 0$ , there exists a time  $\tau$  for which  $|\mathbf{x}_Q(t+\tau) - \mathbf{x}_Q(t)| \leq \varepsilon$  holds for any  $t$ ; (it is a curve that densely fills a ‘tubular’ space);
- *Non-periodic orbit*: orbit  $\mathbf{x}_S(t)$  with no special properties.

The first three are *recurrent states*; the last one a *non-recurrent state*.

## 2. FUNDAMENTALS OF STABILITY THEORY

### ▪ Stability of orbits

Basic idea: An orbit is *stable* if all orbits, originating close to it, remain confined in a small neighborhood; otherwise, it *unstable*. This notion specializes as follows.

#### ➤ Stability of an equilibrium point (Liapunov):

$\mathbf{x}_E$  is *stable* if, for every neighborhood  $\mathcal{U}$  of  $\mathbf{x}_E$ , there exists a neighborhood  $\mathcal{V} \subseteq \mathcal{U}$  of  $\mathbf{x}_E$ , such that an orbit  $\mathbf{x}(t)$  starting in  $\mathcal{V}$  remains in  $\mathcal{U}$  for all  $t \geq 0$ . If, in addition,  $\mathbf{x}(t) \rightarrow \mathbf{x}_E$  as  $t \rightarrow \infty$ , then  $\mathbf{x}_E$  is *asymptotically stable*.

#### ➤ Stability of a general orbit (orbital stability):

$\mathbf{x}_s(t)$  is *orbitally stable* if all orbits, originating from nearby initial points, remain ‘close’ to it, irrespectively of their time-parametrization.

## ■ Quantitative analysis: the variational equation

➤ Orbit  $\mathbf{x}_S(t)$  to be analyzed:

$$\dot{\mathbf{x}}_S(t) = \mathbf{F}(\mathbf{x}_S(t))$$

➤ Perturbed motion:

$$\mathbf{x}(t) = \mathbf{x}_S(t) + \delta\mathbf{x}(t)$$

➤ *Equations for the perturbed motion, Taylor- expanded:*

$$\begin{aligned}\dot{\mathbf{x}}_S(t) + \delta\dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}_S(t) + \delta\mathbf{x}(t)) \\ &= \mathbf{F}(\mathbf{x}_S(t)) + \mathbf{F}_x(\mathbf{x}_S(t))\delta\mathbf{x}(t) + \mathcal{O}(|\delta\mathbf{x}(t)|^2)\end{aligned}$$

i.e.:

$$\delta\dot{\mathbf{x}}(t) = \mathbf{J}_S(t)\delta\mathbf{x}(t) + \mathcal{O}(|\delta\mathbf{x}(t)|^2), \quad \mathbf{J}_S(t) := \mathbf{F}_x(\mathbf{x}_S(t))$$

➤ By linearizing in  $\delta\mathbf{x}(t)$ :

$$\delta\dot{\mathbf{x}}(t) = \mathbf{J}_s(t)\delta\mathbf{x}(t)$$

This is called *the variational equation* (based on  $\mathbf{x}_s(t)$ ); it is generally non-autonomous, since  $\mathbf{x}_s(t)$  depends on  $t$ . Special cases:

○  $\mathbf{x}_s(t)$  is an equilibrium point  $\mathbf{x}_E$ :

$$\delta\dot{\mathbf{x}}(t) = \mathbf{J}_E\delta\mathbf{x}(t), \quad \mathbf{J}_E := \mathbf{F}_x(\mathbf{x}_E)$$

in which  $\mathbf{J}_E$  has *constant coefficients*.

○  $\mathbf{x}_s(t)$  is a periodic orbit  $\mathbf{x}_P(t)$  of period  $T$ :

$$\delta\dot{\mathbf{x}}(t) = \mathbf{J}_P(t)\delta\mathbf{x}(t), \quad \mathbf{J}_P(t) := \mathbf{F}_x(\mathbf{x}_P(t))$$

in which  $\mathbf{J}_P(t) = \mathbf{J}_P(t+T)$  has *periodic coefficients* (Floquet Theory).

We will confine ourselves to equilibrium points.

## ■ Stability of an equilibrium point

We study the stability of an equilibrium point  $\mathbf{x}_E$ .

- Linearized stability:

We first study *linearized stability* of  $\mathbf{x}_E$ , by ignoring the reminder  $O(|\delta\mathbf{x}(t)|^2)$  in the equation for perturbed motion.

➤ Variational equation:

$$\delta\dot{\mathbf{x}}(t) = \mathbf{J}_E \delta\mathbf{x}(t)$$

➤ Eigenvalue problem:

$$(\mathbf{J}_E - \lambda\mathbf{I})\mathbf{u} = \mathbf{0}$$



➤ Discussion:

- if *all the eigenvalues*  $\lambda$  have negative real parts, then  $\delta \mathbf{x}(t) \rightarrow 0$  as  $t \rightarrow \infty$ ; therefore  $\mathbf{x}_E$  is *asymptotically stable*;
- if *at least one eigenvalue*  $\lambda$  has positive real part, then  $\delta \mathbf{x}(t) \rightarrow \infty$  as  $t \rightarrow \infty$ ; therefore  $\mathbf{x}_E$  is *unstable*;
- if *all the eigenvalues*  $\lambda$  have non-positive real parts, and at least one of them has zero real part,  $\mathbf{x}_E$  is a *critical* equilibrium point.

➤ Hyperbolic and non-hyperbolic equilibrium points:

- the equilibria at which all the eigenvalues  $\lambda$  of  $\mathbf{J}_E$  have non-zero real parts are *hyperbolic points*;
- the equilibria at which at least one of the eigenvalues  $\lambda$  has zero real part are *non-hyperbolic points* (also named *critical*).

- Nonlinear stability of *hyperbolic points*:

Since the remainder term  $O(|\delta \mathbf{x}(t)|^2)$  in the nonlinear equation

$$\delta \dot{\mathbf{x}}(t) = \mathbf{J}_S(t) \delta \mathbf{x}(t) + O(|\delta \mathbf{x}(t)|^2)$$

can be made as small as we wish, by selecting a sufficiently small neighborhood of  $\mathbf{x}_E$ , results for linear system apply also to nonlinear system. Therefore:

*A hyperbolic point is asymptotically stable if all the eigenvalues of the Jacobian matrix  $\mathbf{J}_E$  have negative real parts; is unstable if at least one eigenvalue has positive real part.*

- Nonlinear stability of *non-hyperbolic points*:

Nonlinear terms decide the true character of the equilibrium, stable or unstable. Therefore, linear stability analysis fails to give an answer, and a nonlinear analysis is necessary.

➤ Example:


A one-dimensional system admitting a non-hyperbolic equilibrium point  $x_E=0$  is considered:

$$\dot{x} = F(x), \quad x \in \mathbb{R}, \quad F(0) = F_x(0) = 0$$

By letting  $x=x_E+\delta x$  and expanding in series, the equation reads:

$$\delta\dot{x} = \cancel{F(0)} + \cancel{F_x(0)}\delta x + \frac{1}{2}F_{xx}(0)\delta x^2 + \frac{1}{6}F_{xxx}(0)\delta x^3 + O(\delta x^4)$$


The phase-portrait is illustrated in the figure.



### 3. FUNDAMENTS OF BIFURCATION THEORY

Bifurcation theory considers *families of systems* depending on parameters. Its aim is to divide the parameter space in regions in which the system has qualitatively similar behaviors. At the separating boundaries, sudden alteration of the dynamics takes place. They are called *bifurcations*.

Example:



## ■ Parameter-dependent systems

- Autonomous dynamical system:

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t), \boldsymbol{\mu}), \quad \mathbf{x} \in \mathbb{R}^N, \boldsymbol{\mu} \in \mathbb{R}^M$$

where  $\boldsymbol{\mu}$  are parameters.

- Phase-portrait:


Since orbits  $\mathbf{x}_s = \mathbf{x}_s(t, \boldsymbol{\mu})$  depend on parameters, when these latter are (quasi-statically) varied, the whole phase-portrait is modified.

## ■ Structural Stability

A phase-portrait is *robust*, or *structurally stable*, if small perturbations of the vector-field (as  $\varepsilon \mathbf{G}(\mathbf{x}, \boldsymbol{\mu})$ , with  $\varepsilon \ll 1$ ) do not qualitatively change it, but only entails smooth deformations.

- **Note:** Stability and Structural Stability should not be confused. The former refers to a *selected orbit*, and depends on the phase-portraits surrounding it; the latter refer on the *whole phase-portrait*, or, at least, a region of interest (local bifurcation)

➤ Example of robust dynamics:



## ■ Bifurcation: general definition

- A bifurcation is *a qualitative change of dynamics*. It occurs at a bifurcation value  $\mu = \mu_c$  of the parameters at which structural stability is lost. Therefore, the dynamics at  $\mu_c + \delta\mu$ , with  $|\delta\mu|$  arbitrary small, is *topologically inequivalent* from that at  $\mu_c$ .
- **Note:** at a bifurcation, stability of equilibria changes, or the number of equilibria and/or periodic orbits change.

## ■ Bifurcation of equilibrium points:

- If  $\mathbf{x}_E^0$  is hyperbolic, the sign of the real part of the eigenvalues of  $\mathbf{J}_E^0$  does not change under sufficiently small perturbations  $\delta\boldsymbol{\mu}$ , so that the dynamics remain substantially unaltered. *The local phase-portrait at a hyperbolic equilibrium point is structurally stable.*
  - If  $\mathbf{x}_E^0$  is non-hyperbolic, one or more eigenvalues of  $\mathbf{J}_E^0$  have zero real parts. Therefore, arbitrary small perturbations  $\delta\boldsymbol{\mu}$  may lead to eigenvalues of  $\mathbf{J}_E$  with (small) positive or negative real parts, thus strongly changing the dynamics. We conclude that *the local phase-portrait at a non-hyperbolic equilibrium point is structurally unstable.*
- By summarizing:
- Bifurcation of equilibrium occurs at *non-hyperbolic* points.





- Example: saddle-node bifurcation:


Dynamical system:

$$\begin{cases} \dot{x} = \mu + x^2 \\ \dot{y} = -y \end{cases}$$


Equilibrium path



$\mu < 0$



$\mu = 0$



$\mu > 0$

# MULTIPLE BIFURCATIONS FROM A KNOWN PATH

## ▪ Bifurcations from a known path

We consider an autonomous dynamical system depending on parameters:

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t), \boldsymbol{\mu}), \quad \mathbf{x} \in \mathbb{R}^N, \boldsymbol{\mu} \in \mathbb{R}^M$$

We assume to know an equilibrium path (named *fundamental path*)



$\mathbf{x}_E = \mathbf{x}_E(\boldsymbol{\mu})$ ; we want:

- (a) to find the values  $\boldsymbol{\mu}_c$  of the parameters for which a bifurcation takes place (analysis of *critical behavior*);
- (b) to study the dynamics of the nonlinear system for values of  $\boldsymbol{\mu}$  close to  $\boldsymbol{\mu}_c$  (*post-critical behavior*).

For task (a), a linearized analysis is required; for task (b), a nonlinear analysis is necessary.

## ▪ Local form of the equation of motion

It can be assumed that the fundamental path is trivial,  $\tilde{\mathbf{x}}_E = \mathbf{0} \forall \mu$ , since it just requires a change of coordinates:




## ■ Static and dynamic bifurcations

We consider a bifurcation point  $\boldsymbol{\mu} = \boldsymbol{\mu}_c$ , at which the Jacobian matrix

$\mathbf{F}_x^c := \mathbf{F}_x(\mathbf{x}_E(\boldsymbol{\mu}_c), \boldsymbol{\mu}_c)$  admits  $N_c := N_z + 2 N_i$  non-hyperbolic (*critical or central*)

eigenvalues with zero real part, ( $\lambda_k = 0, 0, \dots; \pm i\omega_1, \pm i\omega_2, \dots$ ), the remaining (non-critical) having negative real part (example in Fig:  $N_z=2, N_i=2$ ).



- Transversality condition

We assume that the critical eigenvalues cross the imaginary axis with non-zero velocity (i.e.  $(\partial / \partial \boldsymbol{\mu}) \operatorname{Re}(\lambda_k) \neq \mathbf{0}$  ).

We distinguish:

- *static bifurcation* (also said *divergence bifurcation*), if the critical eigenvalues are all zero,  $\lambda_k = 0, k = 1, 2, \dots, N_z$ ;
- *dynamic bifurcation* (or Hopf bifurcation): if the critical eigenvalues are all purely imaginary,  $\lambda_j = \pm i\omega_j, j = 1, 2, \dots, N_i$ ;
- *static-dynamic bifurcation*: if the critical eigenvalues are either zero and purely imaginary,  $\lambda_k = 0, \lambda_j = \pm i\omega_j, k = 1, 2, \dots, N_z, j = 1, 2, \dots, N_i$ .

If  $N_z + N_i = 1$ , the bifurcation is called *simple*; if  $N_z + N_i > 1$ , the bifurcation is called *multiple*.

## ▪ Resonant dynamic bifurcations

If the critical eigenvalues  $\lambda_j = \pm i\omega_j$  are *rationally dependent*, i.e. if there exist some sets of integer numbers  $k_{rj}$  such that:

$$\sum_{j=1}^{N_c} k_{rj} \omega_j = 0, \quad k_{rj} \in \mathbb{Z}, \quad r = 1, 2, \dots, R$$

then the (multiple) bifurcation is said *resonant*. If no such numbers exist, the bifurcation is said *non-resonant*.

- **Note:** resonance conditions do not play any role in determining if a point is, or not, of bifurcation. However, as it will be shown ahead, the resonances strongly affect the *nonlinear* dynamics.

## ▪ Linear codimension of a bifurcation

In the parameter-space, bifurcations take place on manifolds on which some relations (*constraints*) among the parameters is satisfied, namely:

➤ if the bifurcation is *non-resonant*:

$$\operatorname{Re}(\lambda_k) = 0, \quad k = 1, 2, \dots, N_z + N_i$$

➤ if the bifurcation is *resonant*:


$$\begin{cases} \operatorname{Re}(\lambda_k) = 0, & k = 1, 2, \dots, N_z + N_i \\ \sum_{k=1}^{N_c} k_{rj} \operatorname{Im}(\lambda_k) = 0, & r = 1, 2, \dots, R \end{cases}$$

The number:

$$M := N_z + N_i + R$$

is the (linear) *codimension* of the bifurcation; it is the codimension of the manifold on which the multiple bifurcation occurs.

• Examples of low-codimension bifurcations:




Divergence	Hopf	Double-zero	Hopf-Diverg.	Double-Hopf
$M=1$	$M=1$	$M=2$	$M=2$	$\begin{cases} M = 2 \text{ non-res.} \\ M = 3 \text{ resonant} \end{cases}$
$\lambda_1 = 0$	$\text{Re}(\lambda_1) = 0$	$\begin{cases} \lambda_1 = 0 \\ \lambda_2 = 0 \end{cases}$	$\begin{cases} \text{Re}(\lambda_1) = 0 \\ \lambda_2 = 0 \end{cases}$	$\begin{cases} \text{Re}(\lambda_1) = 0 \\ \text{Re}(\lambda_2) = 0 \end{cases}$ $\omega_2 / \omega_1 = r \in \mathbb{Q}$



## ■ The bifurcation parameters

- A bifurcation point  $\mu_c$  would appear as a rare singularity, if specific systems were considered. A small perturbation  $\delta\mu$  would cancel it.
- In contrast, if a *family of systems* is considered, the bifurcating system *naturally appears* as a member of that family.



- The lowest-dimensional family where to embed the bifurcation has dimension- $M$ , and it is *transversal to the critical manifold*.
- Any perturbation of the transversal manifold changes the bifurcation point, but cannot destroy the bifurcation. The  $M$  parameters of the family are the *bifurcation parameters*.

## ▪ The (nonlinear) bifurcation analysis

To analyze the system dynamics around a bifurcation point  $\mu = \mu_c$ , there are essentially two methods:

- The *Center Manifold Method* (CMM) and *Normal Form Theory* (NFT)
- The *Multiple Scale Method* (MSM) (or equivalent perturbation methods).

The CMM *reduces the dimension* of the system, leading to an equivalent system (bifurcation equations) which describes the asymptotic ( $t \rightarrow \infty$ ) dynamics.

The NFT *reduces the complexity* of the bifurcation equations, giving them the simplest nonlinear form.

The MSM performs both the operations simultaneously. In addition *it filters the fast dynamics* from the bifurcation equations.